Identifying Enrollment Challenges and Discovering Research Opportunities in an Orthopaedic Trauma Consortium: The Value of a "Start-up" Registry

The Major Extremity Trauma Research Consortium (METRC)

www.METRC.org

PURPOSE


The Major Extremity Trauma Research Consortium (METRC) was funded to research major challenges to the successful care of and recovery from severe extremity trauma. To assist in the design of studies that are feasible and adequately powered, a registry was established. At the initiation of the Consortium, many of the centers did not have a registry, and those that did were not collecting data under a uniform protocol. We hypothesized that this registry would provide a good snapshot of the injuries treated by Consortium centers and would: (1) assist in making decisions regarding expansion of the Consortium to more centers; (2) identify recruitment challenges in competing projects; and (3) identify opportunities for additional studies based on the identification of high incident injuries that were not being researched by the Consortium.

ABOUT METRC

The overall goal of the METRC Consortium is to produce the evidence needed to establish treatment guidelines for the optimal care of the wounded warrior and ultimately improve the clinical, functional and quality of life outcomes of both service members and civilians who sustain high energy trauma to the extremities.

Anchored by a Data Coordinating Center at the Johns Hopkins Bloomberg School of Public Health and its Center for Injury Research and Policy, the Consortium includes 22 core Level I civilian trauma centers and 4 core Military Treatment Facilities (MTFs) – with the ability to expand patient recruitment to more than 30 additional satellite trauma centers.

LOCATION OF METRC CENTERS



METHODS

In 2010, an IRB-approved registry was established to assist the Consortium in determining the feasibility of future studies that could address critical research questions with adequate power. All core centers were asked to implement the registry via the REDCap distributed data system and maintain the registry for at least 365 consecutive days.

The registry contains a limited set of data on patients between the ages of 18 and 84 who were admitted with fractures requiring surgery of the upper or lower extremity, pelvis or acetabulum, and foot (calcaneus, talus or crush injuries only).


Excluded from the registry are hip fractures in patients 60 years or older and fractures to the wrist, hand, ankle, clavicle, patella, and the foot other than calcaneus/talus/crush.

Date of disch	To be completed by the STUDY COORDINATOR			Patie	nt Name	Date of admission	
Version 1.24 8/16/2010				1 MRN	r		Date of disabance
Age: Gender: ○ Male ○ Female							/ /
Complete the information below for each fracture meeting registry inclusion criteria. Number fractures sequentic reacture #				S	STUDY COORDINATOR USE ONLY: DO NOT ENTER INTO REDCAP		
Discharge Disposition: O Home O Rehab O Died O Other		F C		100			
Complete the information below for each fracture meeting registry inclusion criteria. Number fractures sequentially at Time of Admission Related to THIS Fracture; skip to meeting a Registry Criteria:							
Complete the information below for each fracture meeting registry inclusion criteria. Number fractures sequential racture #						me O Rehab O Die	d O Other O UNK
Injury 2. Major Amputation Level at Discharge: 3. Initial 4. Gustilk Classifies Cleft No amputation (skip to #4) Hip Disarticulation Surgical Closec Clo					and the second second second second second	teria. Number fractu	res sequentially.
Injury 2. Major Amputation Level at Discharge: 3. Initial 4. Gustilic Cleft No amputation (skip to #4) Hip Disarticulation Surgical Closec	4.414						
Composition						3. Initial	4. Gustilo
O Right O Shoulder Disarticulation O Transfemoral amputation O Grade (pelvis O Elbow Disarticulation O Transtibial only) O Transradial O Wrist Disarticulation O Foot/Ankle Disarticulation O Grade amputation* O Grade *If traumatic amp, this form is complete. 5. AO/OTA Classification: a. Bone, Location and Type B. Bone Loss O None O Bone missing or devascularized but still some contact between proximal and distal fragments O Segmental bone loss ≤ 2 cms O Segmental bone loss ≤ 2 cms O Can be approximated O Cannot be approximated O Cannot be approximated O Cannot be approximated O Artery injury with distal ischemia O Can be approximated O Artery injury with distal ischemia O Can be approximated O Artery injury with distal ischemia O Can be approximated O Artery injury with distal ischemia O Cannot be approximated O Artery injury with distal ischemia O Cannot be approximated O Artery injury with distal ischemia O Cannot be approximated O Artery injury with distal ischemia O Cannot be approximated O Artery injury with distal ischemia O Cannot be approximated O Artery injury with distal ischemia O Cannot be approximated O Artery injury with distal ischemia O Cannot be approximated O Artery injury with distal ischemia O Cannot be approximated O Artery injury with distal ischemia O Cannot be approximated O Artery injury with distal ischemia O Cannot be approximated O Artery injury with distal ischemia O Cannot be approximated O Artery injury with distal ischemia O Cannot be approximated O Artery injury with distal ischemia O Cannot be approximated O Artery injury with distal ischemia o	e:					Amputation Type:	Classification:
b. Bone, Location and Type b. Bone Loss ○ None ○ Bone missing or devascularized but still some contact between proximal and distal fragments ○ Segmental bone loss ≤ 2 cms ○ Segmental bone loss > 2 cms ○ Can be approximated ○ Cannot be approximated ○ Extensive degloving ○ Artery injury with distal ischemia ○ No Cannot be approximated ○ Extensive degloving ○ Noreve Abnormality at Time of Admission Related to THIS Fracture (check all that apply): □ Median Complete ○ Yes ○ No ○ UTA ○ NA □ Ulnar ○ Yes ○ No ○ UTA ○ NA □ Ulnar ○ Yes ○ No ○ UTA ○ NA □ Nerve Adial Complete □ No O UTA ○ NA □ Neadial Complete □ Yes ○ No ○ UTA ○ NA □ Neadial Complete □ Yes ○ No ○ UTA ○ NA □ Neadial Superficial □ Yes ○ No ○ UTA ○ NA □ Neadial Superficial □ Yes ○ No ○ UTA ○ NA □ Neadial Superficial □ Yes ○ No ○ UTA ○ NA □ Neadial PlN □ Yes ○ No ○ UTA ○ NA □ Neadial PlN □ Yes ○ No ○ UTA ○ NA □ Neadial PlN □ Yes ○ No ○ UTA ○ NA □ Neadial PlN □ Yes ○ No ○ UTA ○ NA □ Neadial PlN □ Yes ○ No ○ UTA ○ NA □ Neadial Pln □ Yes ○ No ○ UTA ○ NA □ Neadial Plexus □ Yes ○ No ○ UTA □ Na □ Neadial Ple	Right O NA O (pelvis O only) O	Shoulder D Transhume Elbow Disa Transradial	risarticulation ral articulation	O Transfe O Knee D O Transti	emoral Disarticulation bial	amputation O Traumatic amputation* * If traumatic amp, this	O Closed O Grade I or II O Grade IIIA O Grade IIIB O Grade IIIC
O None O Bone missing or devascularized but still some contact between proximal and distal fragments O Segmental bone loss ≤ 2 cms (If closed fracture, skip to uestion #6) Segmental bone loss > 2 cms O Can be approximated O Cannot be approximated O Extensive degloving O Artery injury with distal ischemia O Artery injury with distal ischemia O No O UTA O NA O UTA O NA C Median Complete O Yes O NO O UTA O NA C Radial Complete O Yes O NO O UTA O NA Radial Complete O Yes O NO O UTA O NA Radial Complete O Yes O NO O UTA O NA Radial PlN O Yes O NO O UTA O NA Radial PlN O Yes O NO O UTA O NA Radial PlN O Yes O NO O UTA O NA C Radial PlN O Yes O NO O UTA O NA C Radial PlN O Yes O NO O UTA O NA C Radial PlN O Yes O NO O UTA O NA C Radial PlN O Yes O NO O UTA O NA C O No muscle in area, no appreciable muscle enecrosis, s muscle fingury with intact muscle function. C Loss of muscle but the muscle remains functional, so muscle injury with intact muscle function. C Loss of muscle but the muscle enecrosis, s muscle injury with intact muscle function. C Loss of muscle but the muscle remains functional, so muscle injury with intact muscle function. C Loss of muscle but the muscle enecrosis, s muscle injury with intact muscle function. C Loss of muscle but the muscle enecrosis, s muscle function. C Loss of muscle but the muscle enecrosis, s muscle function. C Loss of muscle but the muscle enecrosis, so muscle function. C Loss of muscle but the muscle enecrosis, so muscle function. C Loss of muscle but the muscle efect does not approximate. U Loss of muscle but the muscle efect does not approximate. U Loss of muscle but the muscle efect does not approximate. U Loss of muscle but the muscle efect does not approximate. U Loss of muscle but the muscle efect does not approximate. U Loss of muscle but the muscle efect does not approximate. U Loss of muscle but the muscle efect does not approximate. U Loss of muscle but the muscle efect does not approximate. U Loss of muscle but the muscle efect does not approximate. U Loss of musc	O/OTA Clas	sification:					
Segmental bone loss ≤ 2 cms Segmental bone loss ≥ 2 cms Seg	one, Location	and Type				144	4
but still some contact between proximal and distal fragments O Segmental bone loss ≤ 2 cms O Segmental bone loss > 2 cms O Can be approximated O Cannot be approximated O Extensive degloving O No clinically apparent injury with distal ischemia O Median Complete O Yes O No O UTA O NA O Ulnar O Yes O No O UTA O NA C Ulnar O Yes O No O UTA O NA C Ulnar O Yes O No O UTA O NA C Ulnar O Yes O No O UTA O NA C Radial Superficial O Yes O No O UTA O NA C Radial Superficial O Yes O No O UTA O NA C Radial Superficial O Yes O No O UTA O NA C Radial PIN O Yes O No O UTA O NA C Radial PIN O Yes O No O UTA O NA C Radial PIN O Yes O No O UTA O NA C O Yes O No O UTA O NA C O Yes O No O UTA O NA C O Yes O NO O U				wasculariza			
Diaphyseal(42); Simple(A) Proximal and distal fragments Segmental bone loss ≤ 2 cms	"42A" for Til	bia/Fibula,					
Dead muscle, loss of muscle function, partial or compartment excision, complete disruption of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function, partial or compartment excision, complete disruption of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function, partial or compartment excision, complete disruption of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function, partial or compartment excision, complete disruption of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function, partial or compartment excision, complete disruption of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function, partial or component excision, complete disruption of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function, partial or component excision, complete disruption of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function, partial or component excision, complete disruption of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function, partial or component excision, complete disruption of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function, partial or component excision, complete disruption of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function (passed on the function of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle function of a mustendon unit, muscle defect does not approximate. Dead muscle, loss of muscle functio					localized necrosis in the zone of injury that requires excision,		
compartment excision, complete disruption of a must tendon unit, muscle defect does not approximate. Unable to assess I. Skin C. Can be approximated C. Cannot be approximated C. Contamination C. Surface contamination (easily removed not embedded or deep soft tissues) C. Imbedded in bone or deep soft tissues OR high risk environmental conditions (barnyard, fecal, dirty water on the properties of the	(If closed fracture, skip to question #6)						
O Can be approximated Cannot be approximated			O Segmental bone loss > 2 cms		compartment excision, complete disruption of a muscle- tendon unit, muscle defect does not approximate.		
O Cannot be approximated O Extensive degloving O Artery injury without ischemia O Extensive degloving O Artery injury with distal ischemia O Artery injury with distal or deep soft tissues O Imbedded in bone or deep soft tissues OR high risk environmental conditions (barnyard, fecal, dirty water i. Median Complete O Yes O No O UTA O NA D Median AIN O Yes O No O UTA O NA D Median AIN O Yes O No O UTA O NA D Median Complete O Yes O No O UTA O NA D Median Complete O Yes O No O UTA O NA D Median AIN O Yes O No O UTA O NA D Median AIN O Yes O No O UTA O NA D Median AIN O Yes O No O UTA O NA D Median AIN O Yes O No O UTA O NA D Median AIN O Yes O No O UTA			e. Arterial		f. Contamination		
Extensive degloving Schemia Sc	kin	imated			O None or minimal contamination.		
ischemia O Imbedded in bone or deep soft tissues OR high risk environmental conditions (barnyard, fecal, dirty water feed) Nerve Abnormality at Time of Admission Related to THIS Fracture (check all that apply): Median Complete O Yes O No O UTA O NA j. Tibial O Yes O No O UTA Median AIN O Yes O No O UTA O NA k. Common Peroneal O Yes O No O UTA Median Complete O Yes O No O UTA O NA l. Deep Peroneal O Yes O No O UTA Medial Complete O Yes O No O UTA O NA m. Superficial Peroneal O Yes O No O UTA Medial Superficial O Yes O No O UTA O NA n. Medial Plantar O Yes O No O UTA Medial PIN O Yes O NO O UTA O NA n. Medial Plantar O Yes O NO O UTA Medial Pin O Yes O NO O UTA O NA n. Brachial Plexus O Yes O NO O UTA Medial Pin O Yes O NO O UTA O NA n. Brachial Plexus O Yes O NO O UTA Medial Pin O Yes O NO O UTA O NA n. Brachial Plexus O Yes O NO O UTA Medial Pin O Yes O NO O UTA O NA n. Brachial Plexus O Yes O NO O UTA Medial Pin O Yes O NO O UTA O NA n. Brachial Plexus O Yes O NO O UTA Medial Pin O Yes O NO O UTA O NA n. Brachial Plexus O Yes O NO O UTA Medial Pin O Yes O NO O UTA O NA n. Brachial Plexus O Yes O NO O UTA Medial Pin O Yes O NO O UTA O NA n. Brachial Plexus O Yes O NO O UTA Median Complete O Yes O NO O UTA O NA n. Brachial Plexus O Yes O NO O UTA Median Complete O Yes O NO O UTA O NA n. Brachial Plexus O Yes O NO O UTA	Can be approxi						
. Median Complete O Yes O No O UTA O NA j. Tibial O Yes O No O UTA O Median AIN O Yes O No O UTA O NA k. Common Peroneal O Yes O No O UTA O NA l. Deep Peroneal O Yes O No O UTA O NA m. Superficial Peroneal O Yes O No O UTA O NA n. Medial Plantar O Yes O No O UTA O NA n. Medial Plantar O Yes O No O UTA	Can be approxi Cannot be appr	roximated					
Median AIN O Yes O No O UTA O NA k. Common Peroneal O Yes O No O UTA Ulnar O Yes O No O UTA O NA l. Deep Peroneal O Yes O No O UTA Radial Complete O Yes O No O UTA O NA m. Superficial Peroneal O Yes O No O UTA Radial Superficial O Yes O No O UTA O NA n. Medial Plantar O Yes O No O UTA Radial PIN O Yes O No O UTA O NA o. Lateral Plantar O Yes O No O UTA Dobturator O Yes O No O UTA O NA p. Brachial Plexus O Yes O No O UTA Sciatic O Yes O No O UTA O NA q. Lumbar Plexus O Yes O No O UTA	Can be approxi Cannot be appr	roximated			environmentar		
. Ulnar O Yes O No O UTA O NA I. Deep Peroneal O Yes O No O UTA . Radial Complete O Yes O No O UTA O NA II. Deep Peroneal O Yes O No O UTA . Radial Superficial O Yes O No O UTA O NA III. Deep Peroneal O Yes O No O UTA . Radial Superficial O Yes O No O UTA O NA III. Deep Peroneal O Yes O No O UTA III. Deep Peroneal O	Can be approxi Cannot be appr Extensive degle Verve Abnorn	roximated oving nality at Ti	ischemia me of Admission Rela		S Fracture (check all		In OTHER ONLA
Radial Complete O Yes O No O UTA O NA m. Superficial Peroneal O Yes O No O UTA Radial Superficial O Yes O No O UTA O NA n. Medial Plantar O Yes O No O UTA Radial PIN O Yes O No O UTA O NA o. Lateral Plantar O Yes O No O UTA Obturator O Yes O No O UTA O NA p. Brachial Plexus O Yes O No O UTA Sciatic O Yes O No O UTA O NA q. Lumbar Plexus O Yes O No O UTA	Can be approxi Cannot be appr Extensive degle Verve Abnorn Median Compl	roximated oving nality at Ti	me of Admission Rela	A O NA	S Fracture (check all j. Tibial	O Yes O N	
Radial Superficial O Yes O No O UTA O NA n. Medial Plantar O Yes O No O UTA Radial PIN O Yes O No O UTA O NA o. Lateral Plantar O Yes O No O UTA Obturator O Yes O No O UTA O NA p. Brachial Plexus O Yes O No O UTA Sciatic O Yes O No O UTA O NA q. Lumbar Plexus O Yes O No O UTA	Can be approxication of the control	roximated oving nality at Ti	me of Admission Rela O Yes O No O UT O Yes O No O UT	TA O NA	S Fracture (check all j. Tibial k. Common Peror	O Yes O Noneal O Yes O N	No OUTA ONA
. Obturator O Yes O No O UTA O NA p. Brachial Plexus O Yes O No O UTA . Sciatic O Yes O No O UTA O NA q. Lumbar Plexus O Yes O No O UTA	Can be approxication of the control	roximated oving nality at Tilete	me of Admission Rela O Yes O No O UT O Yes O No O UT O Yes O No O UT	TA ONA TA ONA TA ONA	S Fracture (check all j. Tibial k. Common Peror l. Deep Peroneal	O Yes O N O Yes O N O Yes O N	No OUTA ONA No OUTA ONA
. Sciatic O Yes O No O UTA O NA q. Lumbar Plexus O Yes O No O UTA	Can be approxication to the complex of the complex	roximated oving nality at Tilete	me of Admission Rela O Yes O No O UT	TA ONA TA ONA TA ONA TA ONA TA ONA	j. Tibial k. Common Peror l. Deep Peroneal m. Superficial Pero n. Medial Plantar	O Yes O N	No OUTA ONA No OUTA ONA No OUTA ONA No OUTA ONA
	Can be approxicannot be approxication Extensive degle Verve Abnorn Median Comple Median AIN Ulnar Radial Comple Radial Superfication PIN	roximated oving nality at Tilete	ischemia me of Admission Rela O Yes O No O UT	TA O NA	j. Tibial k. Common Peror l. Deep Peroneal m. Superficial Pero n. Medial Plantar o. Lateral Plantar	O Yes O N	No OUTA ONA
	Can be approxicannot be approxication Extensive degle verse Abnorm Median Complement AIN Ulnar Radial Complement Radial Superfication PIN Obturator	roximated oving nality at Tilete	ischemia me of Admission Rela O Yes O No O UT	TA O NA	j. Tibial k. Common Peror l. Deep Peroneal m. Superficial Pero n. Medial Plantar o. Lateral Plantar p. Brachial Plexus	O Yes O N	No OUTA ONA
. Final Definitive Fixation Method Performed or 8. Wound Closure Procedures Used Through or Plann	Can be approxicannot be approxication Extensive degle extensive degle extensive degle extensive degle extensive degle extensive degle extensive ex	roximated oving nality at Tilete	ischemia me of Admission Rela O Yes O No O UT	TA O NA	j. Tibial k. Common Peror l. Deep Peroneal m. Superficial Pero n. Medial Plantar o. Lateral Plantar p. Brachial Plexus	O Yes O N	No OUTA ONA
Planned at Discharge (Check all that apply): Discharge (Check all that apply):	Can be approxicannot be approxicannot be approxication of the proximal complement of the proximal comp	roximated oving nality at Ti lete	ischemia me of Admission Rela O Yes O No O UT	TA O NA	j. Tibial k. Common Peror l. Deep Peroneal m. Superficial Pero n. Medial Plantar o. Lateral Plantar p. Brachial Plexus q. Lumbar Plexus	O Yes O N S O Yes O N O Yes O N	No OUTA ONA
	Can be approxicannot be approxicannot be approxicated to the complete of the c	nality at Ti lete te cial	ischemia me of Admission Rela O Yes O No O UT	TA O NA	j. Tibial k. Common Peror l. Deep Peroneal m. Superficial Pero n. Medial Plantar o. Lateral Plantar p. Brachial Plexus q. Lumbar Plexus b. Wound Closure Pr Discharge (Check all	O Yes O N	No OUTA ONA
O IMN O Primary / DPC O VAC assisted O Plate O Flap – Local muscle O Other (Please specify)	Can be approxication be approxication be approxication be approxication be approxication between the approxication between	nality at Ti lete te cial	ischemia me of Admission Rela O Yes O No O UT	TA O NA	j. Tibial k. Common Peror l. Deep Peroneal m. Superficial Pero n. Medial Plantar o. Lateral Plantar p. Brachial Plexus q. Lumbar Plexus d. Wound Closure Pr Discharge (Check all D NA / Closed Fracture	O Yes O N neal O Yes O N S O Yes O N Tocedures Used Throu that apply): e O STSG	No OUTA ONA
O Other (Please specify) O Other (Please specify) O Flap – Easciocutaneous	Can be approxicannot be approxicannot be approxicated to the complex of the compl	nality at Ti lete te cial	ischemia me of Admission Rela O Yes O No O UT	TA O NA	j. Tibial k. Common Peror l. Deep Peroneal m. Superficial Pero n. Medial Plantar o. Lateral Plantar p. Brachial Plexus q. Lumbar Plexus d. Wound Closure Pr Discharge (Check all D NA / Closed Fracture D Primary / DPC	O Yes O N	No OUTA ONA do OUTA ONA do OUTA ONA

The Table provides annual estimates of the number of 'registry' fractures treated per site and over the 22 core sites who have been contributing to the

registry for at least 90 days.


- Over three quarters (77%) of all fractures are to the lower extremity
- Nearly one quarter (23%) of all fractures are open.
- Of the open fractures, 49% are Gustilo type III (33 % IIIA, 13% IIIB and 3% IIIC).
- An estimated 436 amputations (traumatic and surgical) are registered annually across the 22 sites (54 to the upper limbs and 382 to the lower limbs).



RESULTS

As of August 15, 2012, 22 of the 26 core centers had implemented the registry and had entered cases for greater than 90 days; 18 of these centers have entered cases for 365 consecutive days. An additional 3 centers have just started to enter cases or are poised to begin; one center was unable to get the registry approved at its institution.


- A total of 11,972 patients have been registered across 22 trauma centers
- Data have been entered on 15,043 fractures (1.26 fractures per patient)
- The number of cases registered per month averages 36.4 across centers



CONCLUSIONS

The registry data have added value to the Consortium and its ability to plan for future studies. As can be seen in the Figure ... patients with type III tibia fractures are the subjects of several potentially competing METRC projects—driving the need for the addition of new centers. And, despite a high number of upper extremity, pelvic, and femur fracture patients, these injuries are currently the focus of few METRC studies. This information is being used to design future research.

The Consortium is currently in the process of expanding the registry to include information on nerve injuries.



ARE YOU INTERESTED IN PARTICIPATING IN METRC?

METRC is looking for centers that are interested in becoming involved as satellite centers. If you are interested in applying to become a satellite center, email admin@metrc.org or talk to Dr. Michael Bosse or Dr. Ellen MacKenzie.





